

A universal law for tails of density pdf's in multi-dimensional Burgers turbulence

J. Bec^{1,2}, U. Frisch^{1,2}, K. Khanin³ and B. Villone⁴

¹Observatoire de la Côte d'Azur, Lab. G.D. Cassini
B.P. 4229, F-06304 Nice Cedex 4, FRANCE

²CNLS - Theoretical Division, LANL, Los Alamos, NM 87545, USA

³Isaac Newton Institute for Mathematical Sciences
20 Clarkson Road, Cambridge CB3 0EH, UK

⁴CNR - Istituto di Cosmogeofisica, 10133 Torino, ITALY

Contact e-mail: bec@obs-nice.fr

Abstract

Extending work of E, Khanin, Mazel and Sinai [1] on the one-dimensional Burgers equation, we show that density pdf's have universal power-law tails with exponent $-7/2$. This behavior stems from singularities, other than shocks, whose nature is quite different in one and several dimensions. We briefly discuss the possibility of detecting singularities of Navier–Stokes turbulence using pdf tails.

1 Introduction

In recent years there has been considerable interest in probability density functions (pdf) for Navier–Stokes turbulence. Similar questions can be asked for random solutions of Burgers equation (“burgulence”). We are interested here in the tail behavior of the pdf of the density ρ for solutions to the d -dimensional Burgers equation in the limit of vanishing viscosity ($\nu \rightarrow 0$):

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} = \nu \nabla^2 \mathbf{v}, \quad \mathbf{v} = -\nabla \psi, \quad (1)$$

$$\partial_t \rho + \nabla \cdot (\rho \mathbf{v}) = 0. \quad (2)$$

The initial potential $\psi_0(\mathbf{r}_0)$ and the initial density $\rho_0(\mathbf{r}_0)$ are random functions of the space variable. This problem arises, for example, in the study of large-scale structures in the Universe (see Ref. [2] and references therein). As is well known, the Burgers equation leads to shocks in which the density of matter is infinite. Yet, large but finite densities do not necessarily occur in the neighborhood of shocks. E *et al.* [1] considered a related problem of determining the pdf of the velocity gradient for the one-dimensional Burgers equation with large-scale and white-in-time random forcing. They showed that large negative gradients




Figure 1: Pdf of the velocity gradient at negative values in log-log coordinates, for one-dimensional kicked burgulence. A power law with exponent $-7/2$ is obtained over two decades. Simulation uses a modification of the Fast Legendre Transform algorithm of Ref. [5].

come from “preshocks” (nascent shocks) which contribute a power-law tail with exponent $-7/2$ to the pdf. Preshocks correspond to fast fluid particles catching up for the first time with slow ones. They constitute discrete events in (Eulerian) space time. We have shown that the $-7/2$ law for the pdf of (negative) velocity gradients in one dimension applies also (i) for decaying (unforced) “burgulence” with smooth (i.e. large-scale) random initial conditions [3] and (ii) for the case of a deterministic time and space-periodic force which is a sum of delta functions in time with smooth space dependence [4]. For this “kicked burgulence”, the $-7/2$ law was also obtained numerically (see Fig. 1).

2 “Kurtoparabolic” points and the $-7/2$ law

We turn now to the multi-dimensional problem (1)-(2) with smooth and random initial conditions (for details see Ref. [2]). As is known, the one or multi-dimensional viscous Burgers equation can be solved explicitly by means of the Cole–Hopf transformation. From this, by taking the limit $\nu \rightarrow 0$, a “maximum representation” can be derived for the velocity potential

$$\psi(\mathbf{r}, t) = \max_{\mathbf{r}_0} \left(\psi_0(\mathbf{r}_0) - \frac{|\mathbf{r} - \mathbf{r}_0|^2}{2t} \right). \quad (3)$$

The maximum is achieved at a point at which the gradient of the r.h.s. vanishes, leading to $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}(\mathbf{r}_0, 0)$. From this it follows that $\mathbf{r}_0$ is a Lagrangian

coordinate. The “naive” Lagrangian map $\mathbf{r}_0 \mapsto \mathbf{r}$ given by this relation is however not invertible, except for short times. The requirement that the maximum in (3) is *global* can be recast in geometrical terms by introducing the Lagrangian potential $\varphi(\mathbf{r}_0, t) \equiv t\psi_0(\mathbf{r}_0) - |\mathbf{r}_0|^2/2$ and the “proper” Lagrangian map $\mathbf{r}_0 \mapsto \mathbf{r} \equiv -\nabla\varphi_c(\mathbf{r}_0, t)$, where φ_c is the convex-hull of φ with respect to $\mathbf{r}_0$. For example, in one dimension, the graph of φ_c is obtained by tightly pulling a string over the graph of φ . The graph of φ_c coincides with that of φ at regular points, wherever fluid particles have not yet fallen into shocks. It also contains linear and ruled manifolds associated to the different types of shocks: segments when $d = 1$, triangles and ruled surfaces when $d = 2$, etc. Conservation of mass implies that the density is given, at regular points, by $\rho(\mathbf{r}, t) = \rho_0(\mathbf{r}_0)/J(\mathbf{r}_0, t)$, where J is the Jacobian of the Lagrangian map. (The density is infinite in shocks.) Since the Jacobian is (up to a factor $(-1)^d$) equal to the Hessian of the Lagrangian potential (determinant of the matrix of second space derivatives), it follows that large densities are typically obtained only near parabolic points. However, arbitrarily close to a parabolic point there are generically hyperbolic points where the surface defined by φ crosses its tangent (hyper)plane and which, therefore, do not belong to its convex hull. Yet, there exist in general exceptional “kurtoparabolic” points which are parabolic and belong to the boundary of the set of regular points. Near such points, arbitrarily large densities are obtained. In one dimension, the only kurtoparabolic points are the preshocks which are discrete space-time events in both Eulerian and Lagrangian coordinates. In two and more dimensions, kurtoparabolic points are also born at preshocks but persist in general for a finite time (see Fig. 2). In Eulerian space, they are associated to boundaries of shocks (e.g. end points of shock lines for $d = 2$).




Figure 2: The Lagrangian potential in two dimensions, just after a preshock (left) and in the neighborhood of a kurtoparabolic point (right). Continuous lines: separatrices between the regular part and the ruled surface of the convex hull; dotted lines: vanishing of the Jacobian of the Lagrangian map. A and A' are a pair of kurtoparabolic points born with the shock.

We just indicate general ideas involved in the determination of the density

pdf for random initial conditions (which need not be homogeneous). The determination of the large- ρ tail of the cumulative probability distribution of the density is equivalent to finding the fraction of Eulerian space-time where ρ exceeds a given value. The latter is determined by changing from Eulerian to Lagrangian coordinates and using a suitable normal form (i.e. a Taylor expansion to the relevant order) of the Lagrangian potential near a kurtoparabolic point. The theory is rather different in one dimension and higher dimensions, because kurtoparabolic points are persistent only in the latter case. However, the scaling law for the resulting pdf, namely $\propto \rho^{-7/2}$, is the same in all dimensions. In fact, when $d \geq 2$, two orthogonal spatial directions play the same role as space and time in one dimension.

3 Detecting Navier–Stokes singularities

It is clear that the algebraic tail of the pdf of velocity gradients or of density for burgulence comes from identified singularities. Measurements of pdf's for space or time derivatives of Eulerian velocities for incompressible three-dimensional Navier–Stokes turbulence have not revealed power law tails, but such tails may just have been, so far, “lost in the experimental noise”. There has indeed been considerable speculations about singularities of the Navier–Stokes equations in the inviscid limit. If singularities with divergent gradients are present, they will give power-law tails, at least as intermediate asymptotics when the viscosity is small (the converse is however not true, since statistical effects not related to singularities can also give power laws). The confirmed absence of power laws would probably rule out singularities.

References

- [1] W. E, K. Khanin, A. Mazel and Y.G. Sinai. Probability distribution functions for the random forced Burgers equation. *Phys. Rev. Lett.*, 78:1904–1907, 1997.
- [2] U. Frisch, J. Bec and B. Villone. Singularities and the distribution of density in the Burgers/adhesion model. *Physica D*, 2001, in press. e-print: cond-mat/9912110.
- [3] J. Bec and U. Frisch. Probability distribution functions of derivatives and increments for decaying Burgers turbulence. *Phys. Rev. E*, 61:1395–1402, 2000. e-print: cond-mat/9906047.
- [4] J. Bec, U. Frisch and K. Khanin. Kicked Burgers turbulence. *J. Fluid Mech.*, 416:239–267, 2000. e-print: chao-dyn/9910001.
- [5] A. Noullez and M. Vergassola. A fast algorithm for discrete Legendre transforms. *J. Sci. Comp.*, 9:259–281, 1994.